

Mouse Monoclonal Antibody to

β-Catenin (exon 3)

clone 9G2

biotinylated

Order No.: 0004-100BIOTIN/b-CAT-9G2

Size (μg) 100

Lot No.: 0004S

www.nanotools.de

orders & support:

email: info@nanotools.de

phone: +49-7641-455 670

fax: +49-7641-455 671

03/140307F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref. Cell Line	Epitope	Immunogen
IgG1	human, mouse, dog	ELISA, WB, IP, ICC, IHC (PS)	90 kDa	SW480	exon 3 / α-catenin binding site	recombinant β-catenin

Background and Specificity:

The α-, β- and γ-catenins are cytoplasmic proteins mediating the interaction of Ca²⁺-dependent transmembrane adhesion molecules (cadherins) with the cytoskeletal network. The direct interaction of β-catenin with the cytoplasmic domain of cadherins plays a crucial role for cell-cell adhesion and signal transmission between neighbouring cells. Recent studies indicate that β-catenin may also play a role in tumorigenesis since it forms complexes with the tumor suppressor gene product APC. β-catenin directly interacts and constitutively activates transcription factors of the TCF/LEF gene family. Thus it is proposed that β-catenin plays a dual role not only in the maintenance and regulation of cell-cell interactions but also in the regulation of gene activity.

Mab β-CAT-9G2 specifically interacts with exon 3 (α-catenin-binding site) of β-catenin.

Related Products

mab to b-catenin (N-Term/Exon2)

#0003-100/b-CAT-7D11

mab to b-catenin (Core)

#0005-100/b-CAT-9G10

mab to b-catenin (C-Term/Exon14)

#0002-100/b-CAT-7D8

mab to b-catenin (C-Term)

#0006-100/b-CAT-10H8

mab to dephospho-b-catenin (aa35-50)

#0051-100/b-CAT-7A7

mab to dephospho-b-catenin (aa27-37)

#0052-100/b-CAT-8E4

mab to phospho-b-catenin (pY86)

#0123-100/b-CAT-24E1

mab to phospho-b-catenin (pY654)

#0159-100/b-CAT-1B11

For monoclonal antibodies against alpha-catenin, LEF, TFF3, and E-, M- and N-Cadherin, please refer to our website at www.nanotools.de

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.
Formulation:	liquid; 0.5 mg/ml in PBS/0.09% Na-Azide/PEG and Sucrose.
Reconstitution:	
Stability:	Aliquote and freeze in liquid nitrogen; antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months. Avoid repeated freeze / thaw cycles.
Positive Control:	#0801: Cell lysate from untreated SW480 cells.
Immunoblotting:	0.5 μg/ml for HRPO/ECL detection Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.
Immunoprecipitation:	use at 1 - 10 μg per 10 ⁶ vanadate treated A431 cells
Immunocytochemistry:	use at 0.1-1 μg/ml
ELISA:	use at 0.05 μg/ml

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.