

Mouse Monoclonal Antibody to

SAPK2 δ /p38 δ (N-terminus)

clone 5H7

Order No.: 0053-100/SAPK2d-5H7
Size (μ g) 100
Lot No.: 0053S

www.nanotools.de

orders & support:

email: info@nanotools.de
 phone: +49-7641-455 670
 fax: +49-7641-455 671



05/040613F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human	WB, IHC	42 kDa	A431	N-terminus, interacts with SAPK2 (p38/CSBP)	N-terminal peptide conjugated to KLH

Background and Specificity:

Stress-activated Protein Kinases (SAPKs) are strongly activated in response to adverse stimuli such as heat and osmotic shock, UV light and other DNA-damaging reagents, and inhibitors of protein synthesis. They are also activated strongly in response to agonists that are released or produced under conditions of stress, such as proinflammatory cytokines. Stress-activated Protein Kinase 2 (SAPK2/p38) phosphorylates conserved Ser - Pro or Thr - Pro peptide motifs.

Mab SAPK2d-5H7 specifically interacts with the N-terminus of SAPK2 δ .

Related Products

- mab to SAPK1 α /jnk2 (N-terminus)**
#0190-100/SAPK1a-12C5
- mab to SAPK1 γ /jnk1 (N-terminus)**
#0200-100/SAPK1g-5D10
- mab to SAPK1/2 (T - G/P - pY)**
#0041-100/SAPK1/2-9H8
- mab to SAPK2 α (N-terminus)**
#0034-100/SAPK2a-13D5 (crossreaction with Mxi2)
#0035-100/SAPK2a-20B11 (no crossreaction with Mxi2)
- mab to MAPK 1/2 (pT-E-pY)**
#0012-100/MAPK-12D4
- mab to MAPK 2 (C-terminus)**
#0011-100/MAPK2-6G11
- mab to MAPK 2 (N-terminus)**
#0178-100/MAPK2-6H3
- mab to MAPK 2 (internal sequence)**
#0239-100/MAPK2-12A4
- mab to MAPK7/erk5**
#0223-100/MAPK7/erk5-12F2
- mab to Mxi 2 (N-terminus)**
0046-100/Mxi-2F2

Purification: The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.

Formulation: lyophilized from 1 ml PBS / 0.09 % Na-azide / PEG and Sucrose

Reconstitution: Reconstitute with 1 ml H₂O (15 min, RT).

Stability: For long-term storage, freeze lyophilizate upon arrival (-20°C). Upon reconstitution, aliquote and freeze in liquid nitrogen; reconstituted antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.


Positive Control: #0831: Cell lysate from untreated A431 cells

Immunoblotting: 0.5 μ g/ml for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: ND

Immunocytochemistry: ND

ELISA: use at 0.05 μ g/ml



Detection of endogenous SAPK2 delta

Whole cell lysates of serum starved tumor cells (20.000 cells per lane) were applied to SDS-PAGE and transferred to a PVDF membrane. The immunoblot was probed with mab SAPK2d-5H7 (0.5 μ g/ml) for 1h at RT and developed by ECL (exp. time: 30 sec).

lane 1: MDA-MB 231; lane 2: MDA-MB 468; lane 3: MCF-7; lane 4: T47D; lane 5: SW480; lane 6: SW620

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.