

Mouse Monoclonal Antibody to

Amyloid β A4 (1-42), C-Terminus

clone 8G7

FITC-labelled

Order No.: 0061-100FITC/bA4(42)-8G7

Size (μ g) 100

Lot No.: 0061S

www.nanotools.de

orders & support:

email: info@nanotools.de

phone: +49-7641-455 670

fax: +49-7641-455 671

03/020307F

Isotype	Species	Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human		ICC		none	C-Terminus of Amyloid β A4 (1-42), does not crossreact with β A4 (1-40)	C-terminal peptide conjugated to KLH

Background and Specificity:

The beta-amyloid peptide (beta A4), proteolytically released from the amyloid precursor protein (APP), is the principal component of senile plaques in Alzheimer's disease. Cleavage of APP by alpha-secretase or alternatively by beta-secretase leads to generation and extracellular release of soluble APP peptides, S-APP-alpha and S-APP-beta, respectively, and the retention of corresponding membrane-anchored C-terminal fragments, C83 and C99. Subsequent processing of C83 by gamma-secretase yields P3 peptides. This is the major secretory pathway and is nonamyloidogenic. Alternatively, presenilin/nicastrin-mediated gamma-secretase processing of C99 releases the amyloid beta proteins, amyloid-beta 40 (Abeta40) and amyloid-beta 42 (Abeta42), major components of amyloid plaques, and the cytotoxic C-terminal fragments, gamma-CTF(50), gamma-CTF(57) and gamma-CTF(59).

Mab β A4(42)-8G7 specifically interacts with the C-Terminus of β -Amyloid (1 - 42) and does not crossreact with β -Amyloid (1 - 40).

Related Products

mab to β A4, N-Terminus
#0064-100/bA4N-19H5

mab to β A4, N-Terminus
#0084-100/bA4N-19H11

mab to β A4, N-Terminus
#0195-100/bA4N-7F4

mab to β A4, N-Terminus
#0196-100/bA4N-7F9

mab to β A4, N-Terminus
#0197-100/bA4N-11H3

mab to β A4 (1-40), C-Terminus
#0060-100/bA4(40)-5C3

mab to β A4 (1-40/42), C-Terminus
#0062-100/bA4(40/42)-9F1

mab to β A4 (1-42), C-Terminus
#0061-100/bA4(42)-8G7

mab to β A4 (1-43), C-Terminus
#0095-100/bA4(43)-6G12

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.
Formulation:	liquid; 0.5 mg/ml in PBS/0.09% Na-Azide/PEG and Sucrose
Reconstitution:	
Stability:	Aliquote and freeze in liquid nitrogen; store aliquots frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.
	Avoid repeated freeze / thaw cycles.
Positive Control:	none
Immunoblotting:	Please use unlabelled or biotinylated antibody for this application. See related products.
Immunoprecipitation:	ND
Immunocytochemistry:	use at 1 - 10 μ g/ml
ELISA:	ND

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.