

Mouse Monoclonal Antibody to

Pyk2/CAK β (phospho-Tyr 402)

clone 14F6

Order No.: 0081-100/Pyk2/CAK β -14F6
Size (μ g) 100
Lot No.: 0081S



www.nanotools.de

orders & support:

email: info@nanotools.de
 phone: +49-7641-455 670
 fax: +49-7641-455 671

02/160307F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human, mouse, rat	WB, ELISA	116 kDa	none	phosphotyrosine 402 S D I pY A E I	phosphopeptide conjugated to KLH

Background and Specificity:

Proline-rich tyrosine kinase 2 (Pyk2; also known as cellular adhesion kinase β , CAK β) is a 116 kDa cytoplasmic tyrosine kinase. Pyk2 is rapidly phosphorylated on tyrosine residues in response to activation by G-protein-coupled receptors as well as stress signals. Phosphorylation of Pyk2 leads to activation of stress-activated kinase 1 and 2 (SAPK1 and SAPK2).

Mab Pyk2/CAK β -14F6 specifically interacts with activated Pyk2/CAK β phosphorylated at tyrosine 402. The antibody does not crossreact with the non-phosphorylated form of Pyk2/CAK β nor with unrelated phosphorylation sites.

Related Products

Purification: The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.

Formulation: lyophilized from 1 ml 2 x PBS / 0.09 % Na-azide / PEG and Sucrose.

Reconstitution: Reconstitute with 1 ml H₂O (15 min, RT).

Stability: For long-term storage, freeze lyophilizate upon arrival (-20°C). Upon reconstitution, aliquote and freeze in liquid nitrogen; reconstituted antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.


Positive Control: none

Immunoblotting: 0.5 μ g/ml for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: ND

Immunocytochemistry: ND

ELISA: use at 0.05 μ g/ml



Transactivation of PYK2

Whole cell lysates of serum starved HepG2 tumor cells (20.000 cells per lane) were applied to SDS-PAGE and transferred to PVDF membranes. Immunoblots were probed with mab PYK2-14F6 (0.5 μ g/ ml) for 1h at RT and developed by ECL (exp. time: 30 sec).

lane 1:Co; lane 2: serum; lane 3: EGF; lane 4: H2O2; lane 5: Anisomycin; lane 6: Sorbit; lane 7: Arsen; lane 8: Ceramide;

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.