

Mouse Monoclonal Antibody to

EGFR (phospho-Ser 1047)

clone 1H9

Order No.: 0107-100/EGFR-1H9
Size (µg) 100
Lot No.: 0107S



www.nanotools.de

orders & support:

email: info@nanotools.de
phone: +49-7641-455 670
fax: +49-7641-455 671

04/260207F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human, mouse, dog	WB, ELISA, IP, ICC, Luminex	180 kDa	HepG2	Phosphoserine 1047 R Y S p S D S T	phosphopeptide conjugated to KLH

Background and Specificity:

EGFR/erbB receptors are activated upon binding of EGF and EGF-related growth factors such as TGF alpha, beta-cellulin, Hb-EGF, HRG, or NRG. Binding of these ligands leads to receptor homo- and heterodimerization followed by autophosphorylation and activation of downstream signal transduction pathways (MAPK, PI3K/PKB, and STAT). In addition, EGFR becomes fully activated after phosphorylation of Y845 by src family kinases.

Phosphorylation of Y1045 leads to association with cbl and subsequent receptor degradation.

Phosphorylation of S1047 by CamKinase II leads to attenuation of kinase activity; phosphorylation of T654 (by PKC) and T669 (by MAPK, p38) interferes with receptor endocytosis/recycling.

Mab EGFR-1H9 specifically recognizes EGFR phosphorylated at Serine 1047. The antibody is suitable for Western Blot, Immunocytochemistry and ELISA applications.

Related Products

- mab to EGFR (C-terminus)**
#0007-100/EGFR-13G8
- mab to EGFR (cytoplasmic domain)**
#0168-100/EGFR-10F4
- mab to EGFR (extracellular domain)**
#0209-100/EGFR-20E12
- mab to EGFR (aa 960 - 980)**
#0199-100/EGFR-16F8
- mab to EGFR (N-terminus)**
#0201-100/EGFR-14C8
- mab to phospho-EGFR (pY 845)**
#0116-100/EGFR-12A3
- mab to phospho-EGFR (pY1045)**
#0136-100/EGFR-11C2
- mab to phospho-EGFR (pY1068)**
#0187-100/EGFR-15A2
- mab to phospho-EGFR (pY 1086)**
#0188-100/EGFR-8B8
- mab to phospho-EGFR (pY 1148)**
#0219-100/EGFR-10G12
- mab to phospho-EGFR (pY1173)**
#0008-100/EGFR-9H2
- mab to dephospho-EGFR (Y1173)**
#0009-100/EGFR-20G3
- mab to phospho-EGFR (pT669)**
#0191-100/EGFR-5F10
- mab to phospho-EGFR (pT654)**
#0138-100/EGFR-3F2

For monoclonal antibodies against erbB2, phospho-erbB2, erbB3 and erbB4, as well as against various EGFR downstream targets, please refer to our website at www.nanotools.de

Purification: The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography

Formulation: lyophilized from 1 ml PBS / 0.09 % Na-azide / PEG and Sucrose.

Reconstitution: Reconstitute with 1 ml H₂O (15 min, RT).

Stability: For long-term storage, freeze lyophilizate upon arrival (-20°C). Upon reconstitution, aliquote and freeze in liquid nitrogen; reconstituted antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.


Positive Control: #0812: Cell lysate from pervanadate-treated HepG2 cells

Immunoblotting: 0.5 µg/ml for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: use at 1 - 10 µg per 10⁶ pervanadate-treated A431 cells

Immunocytochemistry use at 0.1-1 µg/ml

ELISA: use at 0.1 µg/ml



EGFR Transactivation

Serum starved HepG2 cells were treated for 15min as indicated. Whole cell lysates were separated by SDS-PAGE (ca 20.000 cells/lane). The immunoblot was probed with mab EGFR-1H9 (0.5 µg/ml) for 1h at RT and developed by ECL (exp. time: 30 sec).

lane 1: control; lane 2: PMA; lane 3: Forskolin; lane 4: LPA; lane 5: Sorbit; lane 6: Anisomycin; lane 7: Ionomycin; lane 8: Taxol

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.