

Mouse Monoclonal Antibody to

Fos (phospho-Ser 374)

clone 34E4

Order No.: 0118-100/Fos-34E4
Size (µg) 100
Lot No.: 0118S



www.nanotools.de

orders & support:

email: info@nanotools.de
 phone: +49-7641-455 670
 fax: +49-7641-455 671

03/080507F

| Isotype | Species Reactivity | Applications | Mol. Weight | Ref.Cell Line | Epitope | Immunogen |
|---------|--------------------|--------------|---------------|---------------|--------------------------------------|----------------------------------|
| IgG1 | human, dog | WB, IHC | ELISA, 50 kDa | HepG2 | phosphoserine 374 S L S p S P T L | phosphopeptide conjugated to KLH |

Background and Specificity:

The immediated early gene product c-Fos is expressed following mitogenic stimulation. c-Fos functions as a sensor for MAPK signal duration. When MAPK activation is transient, MAPK activity declines before accumulation of the c-Fos protein. When MAPK activation is sustained, c-Fos is phosphorylated by MAPK at serine 374. Phosphorylation stabilizes the Fos protein and primes c-Fos for additional phosphorylation at threonine 325.

Mab Fos-34E4 specifically interacts with c-Fos phosphorylated at serine 374. The antibody is suitable for Western blot and ELISA applications.

Related Products

- mab to Fos (N-terminus)**
#0122-100/Fos-8B5
- mab to MAPK 1/2 (pT-E-pY)**
#0012-100/MAPK-12D4
- mab to MAPK 2/erk2 (C-terminus)**
#0011-100/MAPK-6G11
- mab to MAPK 2/erk2 (N-terminus)**
#0178-100/MAPK-6H3
- mab to MAPK 7/erk5**
#0223-100/MAPK7/erk5-12F2
- mab to MEK1 (N-terminus)**
#0186-100/MEK1-10B1
- mab to MEK1 (pS218/222)**
- mab to MEK2 (pS222/226)**
#0174-100/MEK1/2-7E10
- mab to MEK1/2**
#0150-100/MEK1/2-9G3
- mab to MEK2 (N-terminus)**
#0148-100/MEK2-8E8
- mab to MKK3/MAP2K3 (N-terminus)**
#0166-100/MKK3-5F7
- mab to MKK7 (N-terminus)**
#0189-100/MKK7-10F7

Purification: The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography

Formulation: lyophilized from 1 ml 2 x PBS / 0.09 % Na-azide / PEG and Sucrose.

Reconstitution: Reconstitute with 1 ml H₂O (15 min, RT).

Stability: For long-term storage, freeze lyophilizate upon arrival (-20°C). Upon reconstitution, aliquote and freeze in liquid nitrogen; reconstituted antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.


Positive Control: #0813: Cell lysate from EGF-treated HepG2 cells

Immunoblotting: 0.5 µg/ml for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: ND

Immunocytochemistry: ND

ELISA: 0.1 µg/ml (protein ELISA); capture ELISA: ND



fos activation

Serum starved HepG2 cells were incubated with 10 ng/ml EGF for the indicated times. Whole cell lysates were prepared with lysis buffer V19 and separated by SDS-PAGE (ca 20.000 cells/lane). Immunoblots were probed with mab fos-34E4 (0.5 µg/ml) for 1h at room temperature and developed by ECL (exp. time: 30 sec).

lane 1: control; lane 2: 5 min EGF; lane 3: 15 min EGF;
 lane 4: 30 min EGF; lane 5: 1h EGF; lane 6: 2h EGF
 lane 7: 4h EGF; lane 8: 8h EGF

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.