

Mouse Monoclonal Antibody to

SAPK1 α /jnk2 (N-Terminus)

clone 12C5

biotinylated

Order No.: 0190-100BIOTIN/SAPK1 α /jnk2-12C5

Size (μ g) 100

Lot No.: 0190S

www.nanotools.de

orders & support:

email: info@nanotools.de

phone: +49-7641-455 670

fax: +49-7641-455 671

04/080507F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref. Cell Line	Epitope	Immunogen
IgG1	human, mouse, rat, dog	WB, ELISA, IHC	54 kDa	A431	N-terminus	peptide conjugated to hemocyanin

Background and Specificity:

Stress-activated Protein Kinases (SAPKs) are strongly activated in response to adverse stimuli such as heat and osmotic shock, UV light and other DNA-damaging reagents, and inhibitors of protein synthesis. They are also activated strongly in response to agonists that are released or produced under conditions of stress, such as proinflammatory cytokines.

The SAPK1/jnk family consists of 3 isoforms: SAPK1 α /jnk2; SAPK1 β /jnk3; SAPK1 γ /jnk1.

Mab SAPK1 α -12C5 specifically interacts with the N-terminus of SAPK1 α /jnk2 kinases.

Related Products

mab to SAPK1 β /jnk3 (N-terminus)

#0220-100/SAPK1 β /jnk3-4G6

mab to SAPK1 γ /jnk1 (N-terminus)

#0200-100/SAPK1 γ /jnk1-5D10

mab to SAPK1/2 (T - G/P - pY)

#0041-100/SAPK1/2-9H8

mab to SAPK2 α (N-terminus)

#0034-100/SAPK2 α -13D5 (crossreaction with Mxi2)

#0035-100/SAPK2 α -20B11 (no crossreaction with Mxi2)

mab to SAPK2 δ (N-terminus)

#0053-100/SAPK2 δ -5H7

mab to MAPK 1/2 (pT-E-pY)

#0012-100/MAPK-12D4

mab to MAPK 2 (C-terminus)

#0011-100/MAPK2-6G11

mab to MAPK 2 (N-terminus)

#0178-100/MAPK2-6H3

mab to MAPK 2 (internal sequence)

#0239-100/MAPK2-12A4

mab to MAPK7/erk5

#0223-100/MAPK7/erk5-12F2

mab to Mxi 2 (N-terminus)

0046-100/Mxi-2F2

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.
Formulation:	liquid; 0.5 mg/ml in PBS/0.09% Na-Azide/PEG and Sucrose
Reconstitution:	
Stability:	Aliquote and freeze in liquid nitrogen; antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.

Positive Control: #0831: Cell lysate from untreated A431 cells

Immunoblotting: 0.5 μ g/ml for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: ND

Immunocytochemistry: ND

ELISA: use at 0.05 μ g/ml

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.