

Mouse Monoclonal Antibody to

MAPK2/erk2 (internal sequence)

clone 12A4

Biotinylated

Order No.: 0239-100BIOTIN/MAPK2-12A4

Size (µg) 100

Lot No.: 0239S

www.nanotools.de

orders & support:

email: info@nanotools.de

phone: +49-7641-455 670

fax: +49-7641-455 671

01/230307F

Isotype	Species Reactivity	Applications	Mol. Weight	Ref. Cell Line	Epitope	Immunogen
IgG1	human	WB	42 kDa	A431	aa 200 - 250	peptide conjugated to hemocyanin

Background and Specificity:

Extracellular signal/mitogen activated protein kinases (erk/MAPK) are a group of proline-directed serine/threonine kinases that are activated by dual phosphorylation of conserved threonine and tyrosine residues within a characteristic T X Y peptide motif. The mitogen-activated kinases erk1 (MAPK1) and erk2 (MAPK2) acquire full enzymatic activity upon phosphorylation of both threonine and tyrosine residues within the sequence motif T E Y.

Mab MAPK2-12A4 recognizes MAP kinase 2 (erk2) at 42 kDa in Western-blot applications.

Related Products

mab to MEK1 (pS218/222)
mab to MEK2 (pS222/226)
 #0174-100/MEK1/2-7E10
mab to MEK1 (N-terminus)
 #0186-100/MEK1.10B1
mab to MEK1/2
 #0150-100/MEK1/2-9G3
mab to MEK2 (N-terminus)
 #0148-100/MEK2-8E8
mab to MKK3 (N-terminus)
 #0166-100/MKK3-5F7
mab to MKK5 (N-terminus)
 #0224-100/MKK5-14B5
mab to MKK7 (N-terminus)
 #0189-100/MKK7-10F7
mab to MAPK 1/2 (pT-E-pY)
 #0012-100/MAPK-12D4
mab to MAPK 2 (C-terminus)
 #0011-100/MAPK-6G11
mab to MAPK 2 (N-terminus)
 #0178-100/MAPK2-6H3
mab to MAPK7/erk5
 #0223-100/MAPK7/erk5-12F2
mab to Fos (pS374)
 #0118-100/Fos-34E4
mab to Fos (N-terminus)
 #0122-100/Fos-8B5
mab to C-Raf (pS621)
 #0102-100/C-Raf-6B4
mab to C-Raf
 #0120-100/C-Raf-PBB-1

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent ultrafiltration and size exclusion chromatography.
Formulation:	liquid; 0.5 mg/ml in PBS/0.09% Na-Azide/PEG and Sucrose
Reconstitution:	
Stability:	Aliquote and freeze in liquid nitrogen; antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Avoid repeated freeze / thaw cycles.

Positive Control: #0831: Cell lysate from untreated A431 cells

Immunoblotting: 0.5 µg/ml for HRPO/ECL detection
Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

Immunoprecipitation: ND

Immunocytochemistry: ND

ELISA: ND

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.