

## Mouse Monoclonal Antibody to

# LC3 (microtubule-associated protein1 light chain 3B) clone 2G6

**Order No.:** 0260-100/LC3-2G6  
**Size (µg)** 100  
**Lot No.:** 0260S

[www.nanotools.de](http://www.nanotools.de)

**orders & support:**  
email: [info@nanotools.de](mailto:info@nanotools.de)  
phone: +49-7641-455 670  
fax: +49-7641-455 671



06/040808F

| Isotype | Species Reactivity                 | Applications | Mol. Weight                  | Ref.Cell Line | Epitope             | Immunogen                                  |
|---------|------------------------------------|--------------|------------------------------|---------------|---------------------|--|
| IgG1    | human, mouse, rat, monkey, hamster | WB, ICC      | LC3-I: 18kDa<br>LC3-II:16kDa | Neuro 2A      | N-terminus of LC3-B | synthetic peptide conjugated to hemocyanin |

### Background and Specificity:

Autophagy is an alternative process of proteasomal degradation for some long-lived proteins or organelles. Alterations in the autophagic-lysosomal compartment have been linked to neuronal death in many neurodegenerative disorders as well as in transmissible neuronal pathologies (prion diseases). Genetic studies in yeast have shown that Autophagy-defective Gene-8 (Atg-8) represents a specific marker for autophagy. Among the four families of mammalian Atg8-related proteins only LC3 (microtubule-associated protein1 light chain 3) is expressed at sufficient high levels and efficiently recruited to autophagic vesicles in cells and tissues. During autophagy the cytoplasmic form, LC3-I is processed and recruited to autophagosomes, where LC3-II is generated by site specific proteolysis near to the C-terminus. Autophagic vacuoles have been also reported frequently in cardiomyopathies or muscle cells exposed to different experimental settings.

**Mab LC3-2G6** specifically recognizes both forms of endogenous LC3, the cytoplasmic LC3-I (18 kDa) as well as the lipidated form generated during autophagosome and autophagolysosome formation: LC3-II (16 kDa).

**NOTE: We strongly recommend to use PVDF membranes for immunoblot analysis.**

|                        |   |
|------------------------|---|
| <b>Purification:</b>   | The antibody was purified from serum-free cell culture supernatant by subsequent ultrafiltration and size exclusion chromatography. |
| <b>Formulation:</b>    | liquid in PBS/0.09% Na-Azide/PEG and Sucrose/50% Glycerol (1 ml, c = 100 µg/ml)   |
| <b>Reconstitution:</b> |   |
| <b>Stability:</b>      | Aliquote and store at -20°C up to 1 year  |

**Positive Control:** #0911: Cell lysate from untreated Neuro 2A

**Immunoblotting:** 0.5 µg/ml for HRPO/ECL detection  
**Recommended blocking buffer:** Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.

**Immunoprecipitation:** ND

**Immunocytochemistry:** Use at 1- 10 µg/ml (paraformaldehyd/methanol fixation)

**ELISA:** ND

### Related Products

#### mab to LC3

#0231-100/LC-3-5F10

#### mab to LC3

#0261-100/LC3-5H3

#### mab to Beclin

#0240-100/Beclin-12B4

#### Alzheimer Disease

##### mab to βA4 (1-40), C-Terminus

#0060-100/bA4(40)-5C3

##### mab to βA4 (1-42), C-Terminus

#0061-100/bA4(42)-8G7

##### mab to βA4 (1-40/42), C-Terminus

#0062-100/bA4(40/42)-9F1

##### mab to βA4 (1-43), C-Terminus

#0095-100/bA4(43)-6G12

##### mab to βA4, N-Terminus

#0064-100/bA4N-19H5


##### mab to βA4, N-Terminus

#0084-100/bA4N-19H11

##### mab to βA4, N-Terminus

#0197-100/bA4N-11H3

For monoclonal antibodies against PKB/akt, and SAPK/jnk, please refer to our website at [www.nanotools.de](http://www.nanotools.de)



#### Detection of endogenous LC-3

Whole cell lysates of untreated tumor cells were applied to SDS-PAGE and transferred to a PVDF membrane. The immunoblot was probed with mab LC3 - 2G6 (0.5 µg/ml) for 1h at RT and developed by ECL (exp. time: 30 sec).

lane 1: HeLa; lane 2: HepG2; lane 3: HEK 293; lane 4: SH-SY5Y; lane 5: MDCK; lane 6: PC12; lane 7: CMT; lane 8: Neuro2A; lane 9: NIH - 3T3

**All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.**