

Mouse Monoclonal Antibody to

LC3 (microtubule-associated protein1 light chain 3) clone 5H3

Order No.: 0261-100/LC3-5H3
Size (µg) 100
Lot No.: 0261S



www.nanotools.de

orders & support:

email: info@nanotools.de
 phone: +49-7641-455 670
 fax: +49-7641-455 671

05/311013F

Isotype	Species	Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human		WB	LC3I: 18 kDa LC3II: 16 kD	Neuro 2A	internal sequence that is identical in LC3A, LC3B and LC3C.	synthetic peptide conjugated to hemocyanin

Background and Specificity:

Autophagy is an alternative process of proteasomal degradation for some long-lived proteins or organelles. Alterations in the autophagic-lysosomal compartment have been linked to neuronal death in many neurodegenerative disorders as well as in transmissible neuronal pathologies (prion diseases). Genetic studies in yeast have shown that Autophagy-defective Gene-8 (Atg-8) represents a specific marker for autophagy. Among the four families of mammalian Atg8-related proteins only LC3 (microtubule-associated protein1 light chain 3) is expressed at sufficient high levels and efficiently recruited to autophagic vesicles in cells and tissues. During autophagy the cytoplasmic form, LC3-I is processed and recruited to autophagosomes, where LC3-II is generated by site specific proteolysis near to the N-terminus. Autophagic vacuoles have been also reported frequently in cardiomyopathies or muscle cells exposed to different experimental settings.

NOTE: We strongly recommend to use PVDF membranes for immunoblot analysis.

Related Products

- mab to LC3**
#0231-100/LC-3-5F10
- mab to LC3**
#0260-100/LC3-2G6
- mab to Beclin**
#0240-100/Beclin-12B4
- Alzheimer Disease**
- mab to βA4 (1-40), C-Terminus**
#0060-100/bA4(40)-5C3
- mab to βA4 (1-42), C-Terminus**
#0061-100/bA4(42)-8G7
- mab to βA4 (1-40/42), C-Terminus**
#0062-100/bA4(40/42)-9F1
- mab to βA4 (1-43), C-Terminus**
#0095-100/bA4(43)-6G12
- mab to βA4, N-Terminus**
#0064-100/bA4N-19H5
- mab to βA4, N-Terminus**
#0084-100/bA4N-19H11
- mab to βA4, N-Terminus**
#0197-100/bA4N-11H3

For monoclonal antibodies against PKB/akt, and SAPK/jnk, please refer to our website at www.nanotools.de

Purification:	The antibody was purified from serum-free cell culture supernatant by subsequent ultrafiltration and size exclusion chromatography.
Formulation:	liquid in PBS/0.09% Na-Azide/PEG and Sucrose/50% Glycerol (1 ml, c = 100 µg/ml)
Reconstitution:	
Stability:	Aliquote and freeze in liquid nitrogen; antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months. Avoid repeated freeze / thaw cycles.
Positive Control:	#1041/PC3/LC3I and #1042/PC3/LC3II positive controls
Immunoblotting:	0.5 µg/ml for HRPO/ECL detection Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer, e.g. nanoTools product #3031-500/CPPT or #3031-3000/CPPT.
Immunoprecipitation:	ND
Immunocytochemistry:	Use at 1- 10 µg/ml (paraformaldehyd/methanol fixation)
ELISA:	ND

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.